



Differential Equations with Boundary-Value Problems

Dennis G. Zill

Loyola Marymount University





Differential Equations with Boundary-Value Problems, Ninth Edition

Dennis G. Zill

Product Director: Terry Boyle
Product Manager: Kate Schrumpf
Content Developer: Spencer Arritt
Product Assistant: Stephen Previte
Marketing Manager: Ana Albinson
Content Project Manager: Jennifer Risden

Art Director: Vernon Boes

Compositor: MPS Limited

Manufacturing Planner: Becky Cross
Production Service: MPS Limited
Photo Researcher: Lumina Datamatics
Text Researcher: Lumina Datamatics
Copy Editor: Steven Summerlight
Text Designer: Terri Wright
Cover Designer: Terri Wright
Cover Image: STS-114 Crew, NASA

Portions of this text appear in *Advanced Engineering Mathematics*, Sixth Edition, Copyright 2018, Jones & Bartlett Learning, Burlington, MA 01803 and are used with the permission of the publisher.

© 2018, 2013 Cengage Learning

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced or distributed in any form or by any means, except as permitted by U.S. copyright law, without the prior written permission of the copyright owner.

For product information and technology assistance, contact us at Cengage Learning Customer & Sales Support, 1-800-354-9706.

For permission to use material from this text or product, submit all requests online at www.cengage.com/permissions.

Further permissions questions can be e-mailed to permissionrequest@cengage.com.

Library of Congress Control Number: 2016936076

Student Edition:

ISBN: 978-1-305-96579-9 Loose-leaf Edition:

ISBN: 978-1-337-29240-5

Cengage Learning

20 Channel Center Street Boston, MA 02210 USA

Cengage Learning is a leading provider of customized learning solutions with employees residing in nearly 40 different countries and sales in more than 125 countries around the world. Find your local representative at **www.cengage.com.**

 ${\it Cengage Learning products are represented in Canada by Nelson Education, Ltd.}$

To learn more about Cengage Learning Solutions, visit **www.cengage.com.** Purchase any of our products at your local college store or at our preferred online store **www.cengagebrain.com.**

Printed in the United States of America
Print Number: 01 Print Year: 2016

Preface vii



1	Introduction to Dif	ferential Equations	2
---	---------------------	---------------------	---

- **1.1** Definitions and Terminology 3
- **1.2** Initial-Value Problems 15
- 1.3 Differential Equations as Mathematical Models 22

CHAPTER 1 IN REVIEW 34



2 First-Order Differential Equations 36

- **2.1** Solution Curves Without a Solution 37
 - **2.1.1** Direction Fields 37
 - **2.1.2** Autonomous First-Order DEs 39
- **2.2** Separable Equations 47
- **2.3** Linear Equations 55
- **2.4** Exact Equations 64
- **2.5** Solutions by Substitutions 72
- **2.6** A Numerical Method 76

CHAPTER 2 IN REVIEW 81



Modeling with First-Order Differential Equations 84

- 3.1 Linear Models 85
- **3.2** Nonlinear Models 96
- **3.3** Modeling with Systems of First-Order DEs 107

CHAPTER 3 IN REVIEW 114



4 Higher-Order Differential Equations 118

- **4.1** Preliminary Theory—Linear Equations 119
 - **4.1.1** Initial-Value and Boundary-Value Problems 119
 - **4.1.2** Homogeneous Equations 121
 - **4.1.3** Nonhomogeneous Equations 127
- **4.2** Reduction of Order 132
- **4.3** Homogeneous Linear Equations with Constant Coefficients 135
- **4.4** Undetermined Coefficients—Superposition Approach 142
- **4.5** Undetermined Coefficients—Annihilator Approach 152
- **4.6** Variation of Parameters 159
- **4.7** Cauchy-Euler Equations 166

		 4.8.1 Initial-Value Problems 173 4.8.2 Boundary-Value Problems 179 4.9 Solving Systems of Linear DEs by Elimination 183 4.10 Nonlinear Differential Equations 188 CHAPTER 4 IN REVIEW 193 		
Shutterstock	5	Modeling with Higher-Order Differential Equations 196		
Brian A Jackson/Shutterstock		 5.1 Linear Models: Initial-Value Problems 197 5.1.1 Spring/Mass Systems: Free Undamped Motion 197 5.1.2 Spring/Mass Systems: Free Damped Motion 202 5.1.3 Spring/Mass Systems: Driven Motion 204 5.1.4 Series Circuit Analogue 207 		
		5.2 Linear Models: Boundary-Value Problems 2135.3 Nonlinear Models 222		
		CHAPTER 5 IN REVIEW 232		
шо	6	Series Solutions of Linear Equations 236		
Todd Dalton/Shutterstock com		6.1 Review of Power Series 237		
Shutte		6.2 Solutions About Ordinary Points 243		
Dalton		6.3 Solutions About Singular Points 252		
Todd		6.4 Special Functions 262 CHAPTER 6 IN REVIEW 276		
E		The Laplace Transform 278		
ock cor		7.1 Definition of the Laplace Transform 279		
Raimund asy Shutterstock com		7.2 Inverse Transforms and Transforms of Derivatives 286 7.2.1 Inverse Transforms 286		
Rai mun		7.2.2 Transforms of Derivatives 289		
		7.3 Operational Properties I 294 7.3.1 Translation on the <i>s</i> -Axis 295		
		7.3.1 Translation on the <i>s</i> -Axis 295 7.3.2 Translation on the <i>t</i> -Axis 298		
		7.4 Operational Properties II 306		
		7.4.1 Derivatives of a Transform 306		
		7.4.2 Transforms of Integrals 307		
		7.4.3 Transform of a Periodic Function 3137.5 The Dirac Delta Function 318		
		7.6 Systems of Linear Differential Equations 322		
		CHAPTER 7 IN REVIEW 327		
	8	Systems of Linear First-Order Differential		
ideo (ideo		Equations 332		
Pavel L Photo and V//Shutterstock.com		8.1 Preliminary Theory—Linear Systems 333		
I. Phot terstoc		8.2 Homogeneous Linear Systems 340		
Pave		8.2.1 Distinct Real Eigenvalues 3418.2.2 Repeated Eigenvalues 344		
		o.z.z Repeated Eigenvalues 344		

Green's Functions 173

4.8

8.2.3 Complex Eigenvalues 348

8.3	Nonhomogeneous Linear Systems		
	8.3.1 Undetermined Coefficients	355	
	8.3.2 Variation of Parameters 35	7	
8.4			
	CHAPTER 8 IN REVIEW 366		



Numerical Solutions of Ordinary Differential Equations 368

- **9.1** Euler Methods and Error Analysis 369
- **9.2** Runge-Kutta Methods 374
- **9.3** Multistep Methods 378
- **9.4** Higher-Order Equations and Systems 381
- **9.5** Second-Order Boundary-Value Problems 385

CHAPTER 9 IN REVIEW 389



10 Systems of Nonlinear First-Order Differential Equations 390

- **10.1** Autonomous Systems 391
- **10.2** Stability of Linear Systems 397
- **10.3** Linearization and Local Stability 405
- **10.4** Autonomous Systems as Mathematical Models 414

CHAPTER 10 IN REVIEW 422



11 Fourier Series 424

- **11.1** Orthogonal Functions 425
- **11.2** Fourier Series 431
- **11.3** Fourier Cosine and Sine Series 436
- **11.4** Sturm-Liouville Problem 444
- **11.5** Bessel and Legendre Series 451
 - **11.5.1** Fourier-Bessel Series 452
 - **11.5.2** Fourier-Legendre Series 455

CHAPTER 11 IN REVIEW 458



12 Boundary-Value Problems in Rectangular Coordinates 460

- **12.1** Separable Partial Differential Equations 461
- **12.2** Classical PDEs and Boundary-Value Problems 465
- **12.3** Heat Equation 471
- **12.4** Wave Equation 473
- **12.5** Laplace's Equation 479
- **12.6** Nonhomogeneous Boundary-Value Problems 484
- **12.7** Orthogonal Series Expansions 491
- **12.8** Higher-Dimensional Problems 496

CHAPTER 12 IN REVIEW 499







13 Boundary-Value Problems in Other Coordinate Systems 502

- **13.1** Polar Coordinates 503
- **13.2** Polar and Cylindrical Coordinates 508
- **13.3** Spherical Coordinates 515

CHAPTER 13 IN REVIEW 517

14 Integral Transforms 520

- **14.1** Error Function 521
- **14.2** Laplace Transform 522
- **14.3** Fourier Integral 530
- **14.4** Fourier Transforms 536

CHAPTER 14 IN REVIEW 542

15 Numerical Solutions of Partial Differential Equations 544

- **15.1** Laplace's Equation 545
- **15.2** Heat Equation 550
- **15.3** Wave Equation 555

CHAPTER 15 IN REVIEW 559

Appendices

- A Integral-Defined Functions APP-3
- B Matrices APP-11
- C Laplace Transforms APP-29

Answers for Selected Odd-Numbered Problems ANS-1

Index I-1

TO THE STUDENT

Authors of books live with the hope that someone actually *reads* them. Contrary to what you might believe, almost everything in a typical college-level mathematics text is written for you and not the instructor. True, the topics covered in the text are chosen to appeal to instructors because they make the decision on whether to use it in their classes, but everything written in it is aimed directly at you, the student. So I want to encourage you—no, actually I want to tell you—to read this textbook! But do not read this text as you would a novel; you should not read it fast and you should not skip anything. Think of it as a workbook. By this I mean that mathematics should always be read with pencil and paper at the ready because, most likely, you will have to work your way through the examples and the discussion. Before attempting any problems in the section exercise sets, work through all the examples in that section. The examples are constructed to illustrate what I consider the most important aspects of the section, and therefore, reflect the procedures necessary to work most of the problems. When reading an example, copy it down on a piece of paper and do not look at the solution in the book. Try working it, then compare your results against the solution given, and, if necessary resolve any differences. I have tried to include most of the important steps in each example, but if something is not clear you should always try-and here is where the pencil and paper come in again—to fill in the details or missing steps. This may not be easy, but it is part of the learning process. The accumulation of facts followed by the slow assimilation of understanding simply cannot be achieved without a struggle.

Specifically for you, a *Student Resource Manual (SRM)* is available as an optional supplement. In addition to containing solutions of selected problems from the exercises sets, the *SRM* contains hints for solving problems, extra examples, and a review of those areas of algebra and calculus that I feel are particularly important to the successful study of differential equations. Bear in mind you do not have to purchase the *SRM*; you can review the appropriate mathematics from your old precalculus or calculus texts.

In conclusion, I wish you good luck and success. I hope you enjoy the text and the course you are about to embark on—as an undergraduate math major it was one of my favorites because I liked mathematics that connected with the physical world. If you have any comments, or if you find any errors as you read/work your way through the text, or if you come up with a good idea for improving either it or the *SRM*, please feel free to contact me through Cengage Learning:

spencer.arritt@cengage.com.

TO THE INSTRUCTOR

In case you are examining this text for the first time, *Differential Equations with Boundary-Value Problems*, Ninth Edition, can be used for either a one- or two-semester course that covers ordinary and partial differential equations. The shorter version of the text, *A First Course in Differential Equations with Modeling Applications*, Eleventh Edition, is intended for either a one-semester or one-quarter course in ordinary differential equations. This text ends with Chapter 9. For a one-semester course, it is assumed that the students have successfully completed at least two semesters of calculus. Since you are reading this, undoubtedly you have already examined

the table of contents for the topics that are covered. You will not find a "suggested syllabus" in this preface; I will not pretend to be so wise as to tell other teachers what to teach. I feel that there is plenty of material here to choose from and to form a course to your liking. The text strikes a reasonable balance between the analytical, qualitative, and quantitative approaches to the study of differential equations. As far as my "underlying philosophy" goes, it is this: An undergraduate text should be written with the students' understanding kept firmly in mind, which means to me that the material should be presented in a straightforward, readable, and helpful manner, while keeping the level of theory consistent with the notion of a "first course."

For those who are familiar with the previous editions, I would like to mention a few improvements made in this edition.

- Many exercise sets have been updated by the addition of new problems. Some
 of these problems involve new and, I think, interesting mathematical models.
- Additional examples, figures, and remarks have been added to many sections.
- Throughout the text I have given a greater emphasis to the concepts of piecewise-linear differential equations and solutions that involve nonelementary integrals.
- Appendix A, *Integral-Defined Functions*, is new to the text.
- The superposition principle has been added to the discussion in Section 12.4, *Wave Equation*.
- Section 12.6, Nonhomogeneous Boundary-Value Problems, has been rewritten.
- Modified Bessel functions are given a greater emphasis in Section 13.6, Polar and Cylindrical Coordinates.

Student Resources

- Student Resource Manual (SRM), prepared by Warren S. Wright and Roberto Martinez (ISBN 978-1-305-96573-7, accompanies A First Course in Differential Equations with Modeling Applications, Eleventh Edition, and ISBN 978-1-305-96581-2 accompanies Differential Equations with Boundary-Value Problems, Ninth Edition) provides important review material from algebra and calculus, the solution of every third problem in each exercise set (with the exception of the Discussion Problems and Computer Lab Assignments), relevant command syntax for the computer algebra systems Mathematica and Maple, and lists of important concepts, as well as helpful hints on how to start certain problems.
- MindTap for Differential Equations with Boundary-Value Problems, Ninth Edition, is a digital representation of your course that provides you with the tools you need to better manage your limited time, stay organized, and be successful. You can complete assignments whenever and wherever you are ready to learn with course material specially customized for you by your instructor and streamlined in one proven, easy-to-use interface. With an array of study tools, you will get a true understanding of course concepts, achieve better grades, and set the groundwork for your future courses. Learn more at www.cengage.com/mindtap.

Instructor Resources

- *Instructor's Solutions Manual (ISM)*, prepared by Warren S. Wright and Roberto Martinez, provides complete worked-out solutions for all problems in the text. It is available through the Instructor Companion website at **cengage.com.**
- Cengage Learning Testing Powered by Cognero is a flexible online system that allows you to author, edit, and manage test bank content, create multiple test versions in an instant, and deliver tests from your learning management system (LMS), your classroom, or wherever you want. This is available online at www.cengage.com/login.
- Turn the light on with **MindTap for** *Differential Equations with Boundary Value Problems*, Ninth Edition. Through personalized paths of dynamic

assignments and applications, MindTap is a digital learning solution and representation of your course.

The Right Content: With MindTap's carefully curated material, you get the precise content and groundbreaking tools you need for every course you teach.

Personalization: Customize every element of your course—from rearranging the Learning Path to inserting videos and activities.

Improved Workflow: Save time when planning lessons with all of the trusted, most current content you need in one place in MindTap.

Tracking Students' Progress in Real Time: Promote positive outcomes by tracking students in real time and tailoring your course as needed based on the analytics.

Learn more at www.cengage.com/mindtap.

ACKNOWLEDGEMENTS

Compiling a mathematics textbook such as this and making sure that its thousands of symbols and hundreds of equations are accurate is an enormous task, but since I am called "the author," that is my job and responsibility. But many people besides myself have expended enormous amounts of time and energy in working toward its eventual publication. So I would like to take this opportunity to express my sincerest appreciation to everyone—most of them unknown to me—at Cengage Learning and at MPS North America who were involved in the publication of this edition. A special word of thanks goes to Spencer Arritt, Kathryn Schrumpf, Jennifer Risden, Vernon Boes, and Jill Traut for their guidance in the labyrinth of the production process.

Finally, over the years, this text has been improved in a countless number of ways through the suggestions and criticisms of the reviewers. Thus it is fitting to conclude with an acknowledgement of my debt to the following generous people for sharing their expertise and experience.

REVIEWERS OF PAST EDITIONS

William Atherton, Cleveland State University

Philip Bacon, University of Florida

Bruce Bayly, University of Arizona

William H. Beyer, University of Akron

R. G. Bradshaw, Clarkson College

Bernard Brooks, Rochester Institute of Technology

Allen Brown, Wabash Valley College

Dean R. Brown, Youngstown State University

David Buchthal, University of Akron

Nguyen P. Cac, University of Iowa

T. Chow, California State University-Sacramento

Dominic P. Clemence, North Carolina Agricultural and Technical State University

Pasquale Condo, University of Massachusetts-Lowell

Vincent Connolly, Worcester Polytechnic Institute

Philip S. Crooke, Vanderbilt University

Bruce E. Davis, St. Louis Community College at Florissant Valley

Paul W. Davis, Worcester Polytechnic Institute

Richard A. DiDio, La Salle University

James Draper, University of Florida

James M. Edmondson, Santa Barbara City College

John H. Ellison, Grove City College

Raymond Fabec, Louisiana State University

Donna Farrior, University of Tulsa

Robert E. Fennell, Clemson University

W. E. Fitzgibbon, University of Houston

Harvey J. Fletcher, Brigham Young University

Paul J. Gormley, Villanova

Layachi Hadji, University of Alabama

Ruben Hayrapetyan, Kettering University

Terry Herdman, Virginia Polytechnic Institute and State University

Zdzislaw Jackiewicz, Arizona State University

S. K. Jain, Ohio University

Anthony J. John, Southeastern Massachusetts University

David C. Johnson, University of Kentucky-Lexington

Harry L. Johnson, Virginia Polytechnic Institute and State University

Kenneth R. Johnson, North Dakota State University

Joseph Kazimir, East Los Angeles College

J. Keener, University of Arizona

Steve B. Khlief, Tennessee Technological University

Helmut Knaust, The University of Texas at El Paso

C. J. Knickerbocker, Sensis Corporation

Carlon A. Krantz, Kean College of New Jersey

Thomas G. Kudzma, University of Lowell

Alexandra Kurepa, North Carolina A&T State University

G. E. Latta, University of Virginia

Cecelia Laurie, University of Alabama

Mulatu Lemma, Savannah State University

James R. McKinney, California Polytechnic State University

James L. Meek, University of Arkansas

Gary H. Meisters, University of Nebraska-Lincoln

Stephen J. Merrill, Marquette University

Vivien Miller, Mississippi State University

George Moss, Union University

Gerald Mueller, Columbus State Community College

Philip S. Mulry, Colgate University

Martin Nakashima, California State Polytechnic University-Pomona

C. J. Neugebauer, Purdue University

Tyre A. Newton, Washington State University

Brian M. O'Connor, Tennessee Technological University

J. K. Oddson, University of California–Riverside

Carol S. O'Dell, Ohio Northern University

Bruce O'Neill, Milwaukee School of Engineering

A. Peressini, University of Illinois, Urbana-Champaign

J. Perryman, University of Texas at Arlington

Joseph H. Phillips, Sacramento City College

Jacek Polewczak, California State University Northridge

Nancy J. Poxon, California State University-Sacramento

Robert Pruitt, San Jose State University

K. Rager, Metropolitan State College

F. B. Reis, Northeastern University

Brian Rodrigues, California State Polytechnic University

Tom Roe, South Dakota State University

Kimmo I. Rosenthal, Union College

Barbara Shabell, California Polytechnic State University

Seenith Sivasundaram, Embry-Riddle Aeronautical University

Don E. Soash, Hillsborough Community College

F. W. Stallard, Georgia Institute of Technology

Gregory Stein, The Cooper Union

M. B. Tamburro, Georgia Institute of Technology

Patrick Ward, Illinois Central College

Jianping Zhu, University of Akron

Jan Zijlstra, Middle Tennessee State University


Jay Zimmerman, Towson University



Dennis G. Zill Los Angeles, CA

Differential Equations with Boundary-Value Problems

Introduction to Differential Equations



Kevin George/Shutterstock.com

- 1.1 **Definitions and Terminology**
- 1.2 Initial-Value Problems
- 1.3 Differential Equations as Mathematical Models

CHAPTER 1 IN REVIEW

he words differential and equations suggest solving some kind of equation that contains derivatives y', y'', \ldots . Analogous to a course in algebra, in which a good amount of time is spent solving equations such as $x^2 + 5x + 4 = 0$ for the unknown number x, in this course *one* of our tasks will be to solve differential equations such as y'' + 2y' + y = 0 for an unknown function $y = \phi(x)$. As the course unfolds, you will see there is more to the study of differential equations than just mastering methods that mathematicians over past centuries devised to solve them. But first things first. In order to read, study, and be conversant in a specialized subject you have to learn some of the terminology of that discipline. This is the thrust of the first two sections of this chapter. In the last section we briefly examine the link between differential equations and the real world.

1.1 Definitions and Terminology

INTRODUCTION The derivative dy/dx of a function $y = \phi(x)$ is itself another function $\phi'(x)$ found by an appropriate rule. The exponential function $y = e^{0.1x^2}$ is differentiable on the interval $(-\infty, \infty)$ and by the Chain Rule its first derivative is $dy/dx = 0.2xe^{0.1x^2}$. If we replace $e^{0.1x^2}$ on the right-hand side of the last equation by the symbol y, the derivative becomes

$$\frac{dy}{dx} = 0.2xy. (1)$$

Now imagine that a friend of yours simply hands you equation (1)—you have no idea how it was constructed—and asks, *What is the function represented by the symbol y*? You are now face to face with one of the basic problems in this course:

How do you solve an equation such as (1) for the function $y = \phi(x)$?

A DEFINITION The equation that we made up in (1) is called a **differential equation.** Before proceeding any further, let us consider a more precise definition of this concept.

DEFINITION 1.1.1 Differential Equation

An equation containing the derivatives of one or more unknown functions (or dependent variables), with respect to one or more independent variables, is said to be a **differential equation (DE).**

To talk about them, we shall classify differential equations according to **type**, **order**, and **linearity**.

CLASSIFICATION BY TYPE If a differential equation contains only ordinary derivatives of one or more unknown functions with respect to a *single* independent variable, it is said to be an **ordinary differential equation (ODE).** An equation involving partial derivatives of one or more unknown functions of two or more independent variables is called a **partial differential equation (PDE).** Our first example illustrates several of each type of differential equation.

EXAMPLE 1 Types of Differential Equations

(a) The equations

an ODE can contain more than one unknown function

$$\frac{dy}{dx} + 5y = e^x, \quad \frac{d^2y}{dx^2} - \frac{dy}{dx} + 6y = 0, \quad \text{and} \quad \frac{dx}{dt} + \frac{dy}{dt} = 2x + y \tag{2}$$

are examples of ordinary differential equations.

(b) The following equations are partial differential equations:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2} - 2\frac{\partial u}{\partial t}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$
 (3)

^{*}Except for this introductory section, only ordinary differential equations are considered in *A First Course* in *Differential Equations with Modeling Applications*, Eleventh Edition. In that text the word *equation* and the abbreviation DE refer only to ODEs. Partial differential equations or PDEs are considered in the expanded volume *Differential Equations with Boundary-Value Problems*, Ninth Edition.

4

Notice in the third equation that there are two unknown functions and two independent variables in the PDE. This means u and v must be functions of two or more independent variables.

NOTATION Throughout this text ordinary derivatives will be written by using either the **Leibniz notation** dy/dx, d^2y/dx^2 , d^3y/dx^3 , ... or the **prime notation** y', y'', y''', By using the latter notation, the first two differential equations in (2) can be written a little more compactly as $y' + 5y = e^x$ and y'' - y' + 6y = 0. Actually, the prime notation is used to denote only the first three derivatives; the fourth derivative is written y'' instead of y''''. In general, the *n*th derivative of y'' is written y'' or y''. Although less convenient to write and to typeset, the Leibniz notation has an advantage over the prime notation in that it clearly displays both the dependent and independent variables. For example, in the equation

unknown function

or dependent variable

$$\frac{d^2x}{dt^2} + 16x = 0$$
independent variable

it is immediately seen that the symbol x now represents a dependent variable, whereas the independent variable is t. You should also be aware that in physical sciences and engineering, Newton's **dot notation** (derogatorily referred to by some as the "flyspeck" notation) is sometimes used to denote derivatives with respect to time t. Thus the differential equation $d^2s/dt^2 = -32$ becomes $\ddot{s} = -32$. Partial derivatives are often denoted by a **subscript notation** indicating the independent variables. For example, with the subscript notation the second equation in (3) becomes $u_{xx} = u_{tt} - 2u_t$.

CLASSIFICATION BY ORDER The **order of a differential equation** (either ODE or PDE) is the order of the highest derivative in the equation. For example,

second order
$$\sqrt{\frac{d^2y}{dx^2} + 5\left(\frac{dy}{dx}\right)^3 - 4y} = e^x$$

is a second-order ordinary differential equation. In Example 1, the first and third equations in (2) are first-order ODEs, whereas in (3) the first two equations are second-order PDEs. A first-order ordinary differential equation is sometimes written in the **differential form**

$$M(x, y) dx + N(x, y) dy = 0.$$

EXAMPLE 2 Differential Form of a First-Order ODE

If we assume that y is the dependent variable in a first-order ODE, then recall from calculus that the differential dy is defined to be dy = y'dx.

(a) By dividing by the differential dx an alternative form of the equation (y-x)dx + 4x dy = 0 is given by

$$y - x + 4x \frac{dy}{dx} = 0$$
 or equivalently $4x \frac{dy}{dx} + y = x$.

5

(b) By multiplying the differential equation

$$6xy\frac{dy}{dx} + x^2 + y^2 = 0$$

by dx we see that the equation has the alternative differential form

$$(x^2 + y^2) dx + 6xy dy = 0.$$

In symbols we can express an nth-order ordinary differential equation in one dependent variable by the general form

$$F(x, y, y', \dots, y^{(n)}) = 0,$$
 (4)

where F is a real-valued function of n + 2 variables: $x, y, y', \dots, y^{(n)}$. For both practical and theoretical reasons we shall also make the assumption hereafter that it is possible to solve an ordinary differential equation in the form (4) uniquely for the highest derivative $y^{(n)}$ in terms of the remaining n+1 variables. The differential equation

$$\frac{d^{n}y}{dx^{n}} = f(x, y, y', \dots, y^{(n-1)}),$$
 (5)

where f is a real-valued continuous function, is referred to as the **normal form** of (4). Thus when it suits our purposes, we shall use the normal forms

$$\frac{dy}{dx} = f(x, y)$$
 and $\frac{d^2y}{dx^2} = f(x, y, y')$

to represent general first- and second-order ordinary differential equations.

EXAMPLE 3 Normal Form of an ODE

(a) By solving for the derivative dy/dx the normal form of the first-order differential equation

$$4x \frac{dy}{dx} + y = x$$
 is $\frac{dy}{dx} = \frac{x - y}{4x}$.

(b) By solving for the derivative y'' the normal form of the second-order differential equation

$$y'' - y' + 6 = 0$$
 is $y'' = y' - 6y$.

CLASSIFICATION BY LINEARITY An *n*th-order ordinary differential equation (4) is said to be **linear** if *F* is linear in $y, y', \ldots, y^{(n)}$. This means that an *n*th-order ODE is linear when (4) is $a_n(x)y^{(n)} + a_{n-1}(x)y^{(n-1)} + \cdots + a_1(x)y' + a_0(x)y - g(x) = 0$ or

$$a_n(x)\frac{d^n y}{dx^n} + a_{n-1}(x)\frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x)\frac{dy}{dx} + a_0(x)y = g(x).$$
 (6)

Two important special cases of (6) are linear first-order (n = 1) and linear secondorder (n = 2) DEs:

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$
 and $a_2(x) \frac{d^2y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$. (7)

In the additive combination on the left-hand side of equation (6) we see that the characteristic two properties of a linear ODE are as follows:

- The dependent variable y and all its derivatives $y', y'', \dots, y^{(n)}$ are of the first degree, that is, the power of each term involving y is 1.
- The coefficients a_0, a_1, \ldots, a_n of $y, y', \ldots, y^{(n)}$ depend at most on the independent variable x.

A **nonlinear** ordinary differential equation is simply one that is not linear. Nonlinear functions of the dependent variable or its derivatives, such as $\sin y$ or $e^{y'}$, cannot appear in a linear equation.

EXAMPLE 4 Linear and Nonlinear ODEs

(a) The equations

$$(y-x) dx + 4x dy = 0$$
, $y'' - 2y + y = 0$, $x^3 \frac{d^3y}{dx^3} + x \frac{dy}{dx} - 5y = e^x$

are, in turn, *linear* first-, second-, and third-order ordinary differential equations. We have just demonstrated in part (a) of Example 2 that the first equation is linear in the variable y by writing it in the alternative form 4xy' + y = x.

(b) The equations

nonlinear term: nonlinear term: nonlinear term: nonlinear term: power not 1

$$\downarrow \qquad \qquad \downarrow \qquad \qquad$$

are examples of *nonlinear* first-, second-, and fourth-order ordinary differential equations, respectively.

SOLUTIONS As was stated on page 2, one of the goals in this course is to solve, or find solutions of, differential equations. In the next definition we consider the concept of a solution of an ordinary differential equation.

DEFINITION 1.1.2 Solution of an ODE

Any function ϕ , defined on an interval I and possessing at least n derivatives that are continuous on I, which when substituted into an nth-order ordinary differential equation reduces the equation to an identity, is said to be a **solution** of the equation on the interval.

In other words, a solution of an *n*th-order ordinary differential equation (4) is a function ϕ that possesses at least *n* derivatives and for which

$$F(x, \phi(x), \phi'(x), \dots, \phi^{(n)}(x)) = 0$$
 for all x in I .

We say that ϕ satisfies the differential equation on *I*. For our purposes we shall also assume that a solution ϕ is a real-valued function. In our introductory discussion we saw that $y = e^{0.1x^2}$ is a solution of dy/dx = 0.2xy on the interval $(-\infty, \infty)$.

Occasionally, it will be convenient to denote a solution by the alternative symbol y(x).

INTERVAL OF DEFINITION You cannot think *solution* of an ordinary differential equation without simultaneously thinking *interval*. The interval I in Definition 1.1.2 is variously called the **interval of definition**, the **interval of existence**, the **interval of validity**, or the **domain of the solution** and can be an open interval (a, b), a closed interval [a, b], an infinite interval (a, ∞) , and so on.

7

Verify that the indicated function is a solution of the given differential equation on the interval $(-\infty, \infty)$.

(a)
$$\frac{dy}{dx} = xy^{1/2}$$
; $y = \frac{1}{16}x^4$ (b) $y'' - 2y' + y = 0$; $y = xe^x$

SOLUTION One way of verifying that the given function is a solution is to see, after substituting, whether each side of the equation is the same for every *x* in the interval.

(a) From

left-hand side:
$$\frac{dy}{dx} = \frac{1}{16} (4 \cdot x^3) = \frac{1}{4} x^3,$$

right-hand side: $xy^{1/2} = x \cdot \left(\frac{1}{16} x^4\right)^{1/2} = x \cdot \left(\frac{1}{4} x^2\right) = \frac{1}{4} x^3,$

we see that each side of the equation is the same for every real number x. Note that $y^{1/2} = \frac{1}{4} x^2$ is, by definition, the nonnegative square root of $\frac{1}{16} x^4$.


(b) From the derivatives $y' = xe^x + e^x$ and $y'' = xe^x + 2e^x$ we have, for every real number x,

left-hand side:
$$y'' - 2y' + y = (xe^x + 2e^x) - 2(xe^x + e^x) + xe^x = 0,$$


right-hand side: 0.

Note, too, that each differential equation in Example 5 possesses the constant solution y = 0, $-\infty < x < \infty$. A solution of a differential equation that is identically zero on an interval I is said to be a **trivial solution.**

SOLUTION CURVE The graph of a solution ϕ of an ODE is called a **solution curve.** Since ϕ is a differentiable function, it is continuous on its interval I of definition. Thus there may be a difference between the graph of the *function* ϕ and the graph of the *solution* ϕ . Put another way, the domain of the function ϕ need not be the same as the interval I of definition (or domain) of the solution ϕ . Example 6 illustrates the difference.



(a) function $y = 1/x, x \neq 0$



(b) solution y = 1/x, $(0, \infty)$

FIGURE 1.1.1 In Example 6 the function y = 1/x is not the same as the solution y = 1/x

EXAMPLE 6 Function versus Solution

(a) The domain of y = 1/x, considered simply as a *function*, is the set of all real numbers x except 0. When we graph y = 1/x, we plot points in the xy-plane corresponding to a judicious sampling of numbers taken from its domain. The rational function y = 1/x is discontinuous at 0, and its graph, in a neighborhood of the origin, is given in Figure 1.1.1(a). The function y = 1/x is not differentiable at x = 0, since the y-axis (whose equation is x = 0) is a vertical asymptote of the graph.

(b) Now y = 1/x is also a solution of the linear first-order differential equation xy' + y = 0. (Verify.) But when we say that y = 1/x is a *solution* of this DE, we mean that it is a function defined on an interval I on which it is differentiable and satisfies the equation. In other words, y = 1/x is a solution of the DE on *any* interval that does not contain 0, such as (-3, -1), $(\frac{1}{2}, 10)$, $(-\infty, 0)$, or $(0, \infty)$. Because the solution curves defined by y = 1/x for -3 < x < -1 and $\frac{1}{2} < x < 10$ are simply segments, or pieces, of the solution curves defined by y = 1/x for $-\infty < x < 0$ and

 $0 < x < \infty$, respectively, it makes sense to take the interval *I* to be as large as possible. Thus we take *I* to be either $(-\infty, 0)$ or $(0, \infty)$. The solution curve on $(0, \infty)$ is shown in Figure 1.1.1(b).

EXPLICIT AND IMPLICIT SOLUTIONS You should be familiar with the terms *explicit functions* and *implicit functions* from your study of calculus. A solution in which the dependent variable is expressed solely in terms of the independent variable and constants is said to be an **explicit solution**. For our purposes, let us think of an explicit solution as an explicit formula $y = \phi(x)$ that we can manipulate, evaluate, and differentiate using the standard rules. We have just seen in the last two examples that $y = \frac{1}{16}x^4$, $y = xe^x$, and y = 1/x are, in turn, explicit solutions of $dy/dx = xy^{1/2}$, y'' - 2y' + y = 0, and xy' + y = 0. Moreover, the trivial solution y = 0 is an explicit solution of all three equations. When we get down to the business of actually solving some ordinary differential equations, you will see that methods of solution do not always lead directly to an explicit solution $y = \phi(x)$. This is particularly true when we attempt to solve nonlinear first-order differential equations. Often we have to be content with a relation or expression G(x, y) = 0 that defines a solution ϕ implicitly.

DEFINITION 1.1.3 Implicit Solution of an ODE

A relation G(x, y) = 0 is said to be an **implicit solution** of an ordinary differential equation (4) on an interval I, provided that there exists at least one function ϕ that satisfies the relation as well as the differential equation on I.

It is beyond the scope of this course to investigate the conditions under which a relation G(x, y) = 0 defines a differentiable function ϕ . So we shall assume that if the formal implementation of a method of solution leads to a relation G(x, y) = 0, then there exists at least one function ϕ that satisfies both the relation (that is, $G(x, \phi(x)) = 0$) and the differential equation on an interval I. If the implicit solution G(x, y) = 0 is fairly simple, we may be able to solve for y in terms of x and obtain one or more explicit solutions. See (iv) in the *Remarks*.

EXAMPLE 7 Verification of an Implicit Solution


The relation $x^2 + y^2 = 25$ is an implicit solution of the differential equation


$$\frac{dy}{dx} = -\frac{x}{y} \tag{8}$$


on the open interval (-5, 5). By implicit differentiation we obtain

$$\frac{d}{dx}x^2 + \frac{d}{dx}y^2 = \frac{d}{dx}25 \qquad \text{or} \qquad 2x + 2y\frac{dy}{dx} = 0.$$
 (9)

Solving the last equation in (9) for the symbol dy/dx gives (8). Moreover, solving $x^2 + y^2 = 25$ for y in terms of x yields $y = \pm \sqrt{25 - x^2}$. The two functions $y = \phi_1(x) = \sqrt{25 - x^2}$ and $y = \phi_2(x) = -\sqrt{25 - x^2}$ satisfy the relation (that is, $x^2 + \phi_1^2 = 25$ and $x^2 + \phi_2^2 = 25$) and are explicit solutions defined on the interval (-5, 5). The solution curves given in Figures 1.1.2(b) and 1.1.2(c) are segments of the graph of the implicit solution in Figure 1.1.2(a).







- (a) implicit solution
- **(b)** explicit solution
- (c) explicit solution

$$x^2 + y^2 = 25$$

$$v_1 = \sqrt{25 - x^2}, -5 < x < 5$$

$$y_1 = \sqrt{25 - x^2}, -5 < x < 5$$
 $y_2 = -\sqrt{25 - x^2}, -5 < x < 5$

FIGURE 1.1.2 An implicit solution and two explicit solutions of (8) in Example 7

Because the distinction between an explicit solution and an implicit solution should be intuitively clear, we will not belabor the issue by always saying, "Here is an explicit (implicit) solution."

FAMILIES OF SOLUTIONS The study of differential equations is similar to that of integral calculus. When evaluating an antiderivative or indefinite integral in calculus, we use a single constant c of integration. Analogously, we shall see in Chapter 2 that when solving a first-order differential equation F(x, y, y') = 0we usually obtain a solution containing a single constant or parameter c. A solution of F(x, y, y') = 0 containing a constant c is a set of solutions G(x, y, c) = 0called a **one-parameter family of solutions.** When solving an *n*th-order differential equation $F(x, y, y', ..., y^{(n)}) = 0$ we seek an *n*-parameter family of solutions $G(x, y, c_1, c_2, \dots, c_n) = 0$. This means that a single differential equation can possess an infinite number of solutions corresponding to an unlimited number of choices for the parameter(s). A solution of a differential equation that is free of parameters is called a particular solution.

The parameters in a family of solutions such as $G(x, y, c_1, c_2, ..., c_n) = 0$ are arbitrary up to a point. For example, proceeding as in (9) a relation $x^2 + y^2 = c$ formally satisfies (8) for any constant c. However, it is understood that the relation should always make sense in the real number system; thus, if c = -25 we cannot say that $x^2 + y^2 = -25$ is an implicit solution of the differential equation.




FIGURE 1.1.3 Some solutions of DE in part (a) of Example 8

EXAMPLE 8 Particular Solutions

(a) For all real values of c, the one-parameter family $y = cx - x \cos x$ is an explicit solution of the linear first-order equation

$$xy' - y = x^2 \sin x$$

on the interval $(-\infty, \infty)$. (Verify.) Figure 1.1.3 shows the graphs of some particular solutions in this family for various choices of c. The solution $y = -x \cos x$, the blue graph in the figure, is a particular solution corresponding to c = 0.

(b) The two-parameter family $y = c_1 e^x + c_2 x e^x$ is an explicit solution of the linear second-order equation

$$y'' - 2y' + y = 0$$

in part (b) of Example 5. (Verify.) In Figure 1.1.4 we have shown seven of the "double infinity" of solutions in the family. The solution curves in red, green, and blue are the graphs of the particular solutions $y = 5xe^x$ ($c_1 = 0$, $c_2 = 5$), $y = 3e^x$ ($c_1 = 3$, $c_2 = 0$), and $y = 5e^x - 2xe^x$ ($c_1 = 5, c_2 = 2$), respectively.




FIGURE 1.1.4 Some solutions of DE in part (b) of Example 8