


GLOBAL
EDITION



University Calculus

Early Transcendentals

THIRD EDITION

Joel Hass • Maurice D. Weir • George B. Thomas, Jr.

ALWAYS LEARNING

PEARSON

UNIVERSITY CALCULUS EARLY TRANSCENDENTALS

Third Edition
Global Edition

Joel Hass

University of California, Davis

Maurice D. Weir

Naval Postgraduate School

George B. Thomas, Jr.

Massachusetts Institute of Technology

with the assistance of

Christopher Heil

Georgia Institute of Technology

PEARSON

Boston Columbus Indianapolis New York San Francisco
Amsterdam Cape Town Dubai London Madrid Milan Munich Paris Montréal Toronto
Delhi Mexico City São Paulo Sydney Hong Kong Seoul Singapore Taipei Tokyo

Editorial Director: Chris Hoag
Editor in Chief: Deirdre Lynch
Senior Acquisitions Editor: William Hoffman
Editorial Assistant: Salena Casha
Program Manager: Danielle S. Miller
Project Manager: Rachel S. Reeve
Program Management Team Lead: Marianne Stepanian
Project Management Team Lead: Christina Lepre
Assistant Acquisitions Editor, Global Edition: Aditee Agarwal
Project Editor, Global Edition: Amrita Naskar
Manager, Media Production, Global Edition: Vikram Kumar
Senior Manufacturing Controller, Production, Global Edition: Trudy Kimber
Media Producer: Stephanie Green
TestGen Content Manager: John Flanagan
MathXL Content Manager: Kristina Evans
Marketing Manager: Jeff Weidenaar
Marketing Assistant: Brooke Smith
Senior Author Support/Technology Specialist: Joe Vetere
Rights and Permissions Project Manager: Diahanne Lucas
Procurement Specialist: Carol Melville
Associate Director of Design: Andrea Nix
Program Design Lead: Beth Paquin
Illustrations: Karen Hartpence, Cenveo® Publisher Services
Cover Designer: Lumina Datamatics Ltd.

Cover Image: Teun van den Dries/Shutterstock

Acknowledgments of third party content appear on page 14, which constitutes an extension of this copyright page.

Pearson Education Limited
Edinburgh Gate
Harlow
Essex CM20 2JE
England

and Associated Companies throughout the world

Visit us on the World Wide Web at:
www.pearsonglobaleditions.com

© Pearson Education Limited 2016

The rights of Joel Hass, Maurice D. Weir, and George B. Thomas, Jr. to be identified as the authors of this work have been asserted by them in accordance with the Copyright, Designs and Patents Act 1988.

Authorized adaptation from the United States edition, entitled University Calculus: Early Transcendentals, 3rd edition, ISBN 978-0-321-99958-0, by Joel Hass, Maurice D. Weir, and George B. Thomas, Jr., published by Pearson Education © 2016.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without either the prior written permission of the publisher or a license permitting restricted copying in the United Kingdom issued by the Copyright Licensing Agency Ltd, Saffron House, 6–10 Kirby Street, London EC 1N 8TS.

All trademarks used herein are the property of their respective owners. The use of any trademark in this text does not vest in the author or publisher any trademark ownership rights in such trademarks, nor does the use of such trademarks imply any affiliation with or endorsement of this book by such owners.

PEARSON, ALWAYS LEARNING, MyMathLab and TestGen are exclusive trademarks in the U.S. and/or other countries owned by Pearson Education, Inc. or its affiliates.

Unless otherwise indicated herein, any third-party trademarks that may appear in this work are the property of their respective owners and any references to third-party trademarks, logos or other trade dress are for demonstrative or descriptive purposes only. Such references are not intended to imply any sponsorship, endorsement, authorization, or promotion of Pearson's products by the owners of such marks, or any relationship between the owner and Pearson Education, Inc. or its affiliates, authors, licensees or distributors.

ISBN 10: 1-292-10403-1
ISBN 13: 978-1-292-10403-4

British Library Cataloguing-in-Publication Data
A catalogue record for this book is available from the British Library

10 9 8 7 6 5 4 3 2 1

Typeset by Cenveo Publisher Services

Printed and bound by Vivar in Malaysia

Contents

Preface 9

Credits 14

1

Functions 15

- 1.1 Functions and Their Graphs 15
- 1.2 Combining Functions; Shifting and Scaling Graphs 28
- 1.3 Trigonometric Functions 35
- 1.4 Graphing with Software 43
- 1.5 Exponential Functions 47
- 1.6 Inverse Functions and Logarithms 52

2

Limits and Continuity 65

- 2.1 Rates of Change and Tangents to Curves 65
- 2.2 Limit of a Function and Limit Laws 72
- 2.3 The Precise Definition of a Limit 83
- 2.4 One-Sided Limits 92
- 2.5 Continuity 99
- 2.6 Limits Involving Infinity; Asymptotes of Graphs 110
- [Questions to Guide Your Review](#) 124
- [Practice Exercises](#) 124
- [Additional and Advanced Exercises](#) 126

3

Derivatives 129

- 3.1 Tangents and the Derivative at a Point 129
- 3.2 The Derivative as a Function 133
- 3.3 Differentiation Rules 142
- 3.4 The Derivative as a Rate of Change 152
- 3.5 Derivatives of Trigonometric Functions 161
- 3.6 The Chain Rule 167
- 3.7 Implicit Differentiation 175
- 3.8 Derivatives of Inverse Functions and Logarithms 180
- 3.9 Inverse Trigonometric Functions 190
- 3.10 Related Rates 196
- 3.11 Linearization and Differentials 204
- [Questions to Guide Your Review](#) 215
- [Practice Exercises](#) 216
- [Additional and Advanced Exercises](#) 220

4	Applications of Derivatives	223
4.1	Extreme Values of Functions	223
4.2	The Mean Value Theorem	231
4.3	Monotonic Functions and the First Derivative Test	239
4.4	Concavity and Curve Sketching	244
4.5	Indeterminate Forms and L'Hôpital's Rule	255
4.6	Applied Optimization	264
4.7	Newton's Method	275
4.8	Antiderivatives	279
	Questions to Guide Your Review	289
	Practice Exercises	290
	Additional and Advanced Exercises	294
5	Integrals	297
5.1	Area and Estimating with Finite Sums	297
5.2	Sigma Notation and Limits of Finite Sums	307
5.3	The Definite Integral	314
5.4	The Fundamental Theorem of Calculus	326
5.5	Indefinite Integrals and the Substitution Method	337
5.6	Definite Integral Substitutions and the Area Between Curves	345
	Questions to Guide Your Review	355
	Practice Exercises	355
	Additional and Advanced Exercises	358
6	Applications of Definite Integrals	361
6.1	Volumes Using Cross-Sections	361
6.2	Volumes Using Cylindrical Shells	372
6.3	Arc Length	380
6.4	Areas of Surfaces of Revolution	386
6.5	Work	390
6.6	Moments and Centers of Mass	396
	Questions to Guide Your Review	404
	Practice Exercises	404
	Additional and Advanced Exercises	406
7	Integrals and Transcendental Functions	407
7.1	The Logarithm Defined as an Integral	407
7.2	Exponential Change and Separable Differential Equations	417
7.3	Hyperbolic Functions	426
	Questions to Guide Your Review	434
	Practice Exercises	434
	Additional and Advanced Exercises	435

- ## 8 Techniques of Integration 436
- 8.1 Integration by Parts 437
 - 8.2 Trigonometric Integrals 443
 - 8.3 Trigonometric Substitutions 449
 - 8.4 Integration of Rational Functions by Partial Fractions 454
 - 8.5 Integral Tables and Computer Algebra Systems 461
 - 8.6 Numerical Integration 466
 - 8.7 Improper Integrals 476
 - [Questions to Guide Your Review](#) 487
 - [Practice Exercises](#) 487
 - [Additional and Advanced Exercises](#) 490
- ## 9 Infinite Sequences and Series 492
- 9.1 Sequences 492
 - 9.2 Infinite Series 504
 - 9.3 The Integral Test 513
 - 9.4 Comparison Tests 520
 - 9.5 Absolute Convergence; The Ratio and Root Tests 524
 - 9.6 Alternating Series and Conditional Convergence 530
 - 9.7 Power Series 536
 - 9.8 Taylor and Maclaurin Series 546
 - 9.9 Convergence of Taylor Series 551
 - 9.10 The Binomial Series and Applications of Taylor Series 557
 - [Questions to Guide Your Review](#) 565
 - [Practice Exercises](#) 566
 - [Additional and Advanced Exercises](#) 568
- ## 10 Parametric Equations and Polar Coordinates 571
- 10.1 Parametrizations of Plane Curves 571
 - 10.2 Calculus with Parametric Curves 578
 - 10.3 Polar Coordinates 588
 - 10.4 Graphing Polar Coordinate Equations 592
 - 10.5 Areas and Lengths in Polar Coordinates 595
 - 10.6 Conics in Polar Coordinates 600
 - [Questions to Guide Your Review](#) 606
 - [Practice Exercises](#) 607
 - [Additional and Advanced Exercises](#) 608
- ## 11 Vectors and the Geometry of Space 610
- 11.1 Three-Dimensional Coordinate Systems 610
 - 11.2 Vectors 615
 - 11.3 The Dot Product 624
 - 11.4 The Cross Product 632

- 11.5 Lines and Planes in Space 638
- 11.6 Cylinders and Quadric Surfaces 646
 - [Questions to Guide Your Review](#) 651
 - [Practice Exercises](#) 652
 - [Additional and Advanced Exercises](#) 654

12 Vector-Valued Functions and Motion in Space 656

- 12.1 Curves in Space and Their Tangents 656
- 12.2 Integrals of Vector Functions; Projectile Motion 664
- 12.3 Arc Length in Space 670
- 12.4 Curvature and Normal Vectors of a Curve 675
- 12.5 Tangential and Normal Components of Acceleration 681
- 12.6 Velocity and Acceleration in Polar Coordinates 683
 - [Questions to Guide Your Review](#) 687
 - [Practice Exercises](#) 687
 - [Additional and Advanced Exercises](#) 689

13 Partial Derivatives 690

- 13.1 Functions of Several Variables 690
- 13.2 Limits and Continuity in Higher Dimensions 698
- 13.3 Partial Derivatives 707
- 13.4 The Chain Rule 718
- 13.5 Directional Derivatives and Gradient Vectors 727
- 13.6 Tangent Planes and Differentials 735
- 13.7 Extreme Values and Saddle Points 744
- 13.8 Lagrange Multipliers 753
 - [Questions to Guide Your Review](#) 762
 - [Practice Exercises](#) 763
 - [Additional and Advanced Exercises](#) 766

14 Multiple Integrals 769

- 14.1 Double and Iterated Integrals over Rectangles 769
- 14.2 Double Integrals over General Regions 774
- 14.3 Area by Double Integration 783
- 14.4 Double Integrals in Polar Form 787
- 14.5 Triple Integrals in Rectangular Coordinates 793
- 14.6 Moments and Centers of Mass 802
- 14.7 Triple Integrals in Cylindrical and Spherical Coordinates 809
- 14.8 Substitutions in Multiple Integrals 820
 - [Questions to Guide Your Review](#) 830
 - [Practice Exercises](#) 830
 - [Additional and Advanced Exercises](#) 833

15	Integrals and Vector Fields	835
15.1	Line Integrals	835
15.2	Vector Fields and Line Integrals: Work, Circulation, and Flux	842
15.3	Path Independence, Conservative Fields, and Potential Functions	854
15.4	Green's Theorem in the Plane	865
15.5	Surfaces and Area	877
15.6	Surface Integrals	888
15.7	Stokes' Theorem	899
15.8	The Divergence Theorem and a Unified Theory	911
	Questions to Guide Your Review	922
	Practice Exercises	922
	Additional and Advanced Exercises	925

16	First-Order Differential Equations	online
16.1	Solutions, Slope Fields, and Euler's Method	
16.2	First-Order Linear Equations	
16.3	Applications	
16.4	Graphical Solutions of Autonomous Equations	
16.5	Systems of Equations and Phase Planes	

17	Second-Order Differential Equations	online
17.1	Second-Order Linear Equations	
17.2	Nonhomogeneous Linear Equations	
17.3	Applications	
17.4	Euler Equations	
17.5	Power Series Solutions	

Appendices 927

A.1	Real Numbers and the Real Line	927
A.2	Mathematical Induction	932
A.3	Lines and Circles	936
A.4	Conic Sections	942
A.5	Proofs of Limit Theorems	950
A.6	Commonly Occurring Limits	953
A.7	Theory of the Real Numbers	954
A.8	Complex Numbers	957
A.9	The Distributive Law for Vector Cross Products	965
A.10	The Mixed Derivative Theorem and the Increment Theorem	967

Answers to Odd-Numbered Exercises 971

Index 1035

A Brief Table of Integrals 1051

This page intentionally left blank

Preface

This third edition of *University Calculus* provides a streamlined treatment of the material in a standard three-semester or four-quarter course taught at the university level. As the title suggests, the book aims to go beyond what many students may have seen at the high school level. By emphasizing rigor and mathematical precision, supported with examples and exercises, this book encourages students to think more clearly than if they were using rote procedures.

Generalization drives the development of calculus and is pervasive in this book. Slopes of lines generalize to slopes of curves, lengths of line segments to lengths of curves, areas and volumes of regular geometric figures to areas and volumes of shapes with curved boundaries, rational exponents to irrational ones, and finite sums to series. Plane analytic geometry generalizes to the geometry of space, and single variable calculus to the calculus of many variables. Generalization weaves together the many threads of calculus into an elegant tapestry that is rich in ideas and their applications.

Mastering this beautiful subject is its own reward, but the real gift of mastery is the ability to think through problems clearly—distinguishing between what is known and what is assumed, and using a logical sequence of steps to reach a solution. We intend this book to capture the richness and powerful applicability of calculus, and to support student thinking and understanding for mastery of the material.

New to this Edition

In this new edition, we have followed the basic structure of earlier editions. Taking into account helpful suggestions from readers and users of previous editions, we continued to improve clarity and readability. We also made the following improvements:

- Updated and added numerous exercises throughout, with emphasis on the mid-level and more in the life science areas
- Reworked many figures and added new ones
- Moved the discussion of conditional convergence to follow the Alternating Series Test
- Enhanced the discussion defining differentiability for functions of several variables with more emphasis on linearization
- Showed that the derivative along a path generalizes the single-variable chain rule
- Added more geometric insight into the idea of multiple integrals and the meaning of the Jacobian in substitutions for their evaluations
- Developed surface integrals of vector fields as generalizations of line integrals
- Extended and clarified the discussion of the curl and divergence, and added new figures to help visualize their meanings

Continuing Features

RIGOR The level of rigor is consistent with that of earlier editions. We continue to distinguish between formal and informal discussions and to point out their differences. We think starting with a more intuitive, less formal, approach helps students understand a new or difficult concept so they can then appreciate its full mathematical precision and outcomes. We pay attention to defining ideas carefully and to proving theorems appropriate for calculus students, while mentioning deeper or subtler issues they would study in a more advanced course. Our organization and distinctions between informal and formal discussions give the instructor a degree of flexibility in the amount and depth of coverage of the various topics. For example, while we do not prove the Intermediate Value Theorem or the Extreme Value Theorem for continuous functions on $a \leq x \leq b$, we do state these theorems precisely, illustrate their meanings in numerous examples, and use them to prove other important results. Furthermore, for those instructors who desire greater depth of coverage, in Appendix 7 we discuss the reliance of these theorems on the completeness of the real numbers.

WRITING EXERCISES Writing exercises placed throughout the text ask students to explore and explain a variety of calculus concepts and applications. In addition, the end of each chapter contains a list of questions to help students review and summarize what they have learned. Many of these exercises make good writing assignments.

END-OF-CHAPTER REVIEWS In addition to problems appearing after each section, each chapter culminates with review questions, practice exercises covering the entire chapter, and a series of Additional and Advanced Exercises providing more challenging or synthesizing problems.

WRITING AND APPLICATIONS As always, this text continues to be easy to read, conversational, and mathematically rich. Each new topic is motivated by clear, easy-to-understand examples and is then reinforced by its application to real-world problems of immediate interest to students. A hallmark of this book has been the application of calculus to science and engineering. These applied problems have been updated, improved, and extended continually over the past several editions.

TECHNOLOGY In a course using the text, technology can be incorporated at the discretion of the instructor. Each section contains exercises requiring the use of technology; these are marked with a “T” if suitable for calculator or computer use, or they are labeled “Computer Explorations” if a computer algebra system (CAS, such as Maple or Mathematica) is required.

Additional Resources

INSTRUCTOR'S SOLUTIONS MANUAL (download only)

Single Variable Calculus (Chapters 1–10)

Multivariable Calculus (Chapters 9–15)

The *Instructor's Solutions Manual* contains complete worked-out solutions to all of the exercises. These files are available to qualified instructors through the Pearson Instructor Resource Center, www.pearsonglobaleditions.com/Hass, and MyMathLab.

STUDENT'S SOLUTIONS MANUAL

The *Student's Solutions Manual* is designed for the student and contains carefully worked-out solutions to all the odd-numbered exercises.

JUST-IN-TIME ALGEBRA AND TRIGONOMETRY FOR EARLY TRANSCENDENTALS CALCULUS, Fourth Edition

ISBN 0-321-67103-1 | 978-0-321-67103-5

Sharp algebra and trigonometry skills are critical to mastering calculus, and *Just-in-Time Algebra and Trigonometry for Early Transcendentals Calculus* by Guntram Mueller and Ronald I. Brent is designed to bolster these skills while students study calculus. As students make their way through calculus, this text is with them every step of the way, showing the necessary algebra or trigonometry topics and pointing out potential problem spots. The easy-to-use table of contents has algebra and trigonometry topics arranged in the order in which students will need them as they study calculus.

Technology Resource Manuals (download only)

Maple Manual by Marie Vanisko, Carroll College

Mathematica Manual by Marie Vanisko, Carroll College

TI-Graphing Calculator Manual by Elaine McDonald-Newman, Sonoma State University

These manuals cover Maple 17, Mathematica 8, and the TI-83 Plus/TI-84 Plus and TI-89, respectively. Each manual provides detailed guidance for integrating a specific software package or graphing calculator throughout the course, including syntax and commands.

These manuals are available to qualified instructors through the *University Calculus: Early Transcendentals* Web site, www.pearsonglobaleditions.com/Hass, and MyMathLab.

WEB SITE www.pearsonglobaleditions.com/Hass

The *University Calculus: Early Transcendentals* Web site contains the chapters on First-Order Differential Equations and Second-Order Differential Equations, including odd-numbered answers, and provides the expanded historical biographies and essays referenced in the text. The Technology Resource Manuals and the **Technology Application Projects**, which can be used as projects by individual students or groups of students, are also available.

MyMathLab® Online Course (access code required)

MyMathLab from Pearson is the world's leading online resource in mathematics, integrating interactive homework, assessment, and media in a flexible, easy-to-use format. It provides engaging experiences that personalize, stimulate, and measure learning for each student. And, it comes from an experienced partner with educational expertise and an eye on the future.

To learn more about how MyMathLab combines proven learning applications with powerful assessment, visit www.mymathlab.com or contact your Pearson representative.

Video Lectures with Optional Captioning

The Video Lectures with Optional Captioning feature an engaging team of mathematics instructors who present comprehensive coverage of topics in the text. The lecturers' presentations include examples and exercises from the text and support an approach that emphasizes visualization and problem solving. Available only through MyMathLab.

TestGen®

TestGen® (www.pearsoned.com/testgen) enables instructors to build, edit, print, and administer tests using a computerized bank of questions developed to cover all the objectives of the text. TestGen is algorithmically based, allowing instructors to create multiple but equivalent versions of the same question or test with the click of a button. Instructors can also modify test bank questions or add new questions. The software and test bank are available for download from Pearson Education's online catalog.

PowerPoint® Lecture Slides

These classroom presentation slides are geared specifically to the sequence and philosophy of *University Calculus: Early Transcendentals*. Key graphics from the book are included to help bring the concepts alive in the classroom. These files are available to qualified instructors through the Pearson Instructor Resource Center, www.pearsonglobaleditions.com/Hass, and MyMathLab.

Acknowledgments

We wish to express our gratitude to the reviewers of this and previous editions, who provided such invaluable insight and comment.

Harry Allen, <i>Ohio State University</i>	Hideaki Kaneko, <i>Old Dominion University</i>
Edoh Amiran, <i>Western Washington University</i>	Przemo Kranz, <i>University of Mississippi</i>
Anthony Bedenikovic, <i>Bradley University</i>	John Kroll, <i>Old Dominion University</i>
Robert A. Beezer, <i>University of Puget Sound</i>	Krystyna Kuperberg, <i>Auburn University</i>
Przemyslaw Bogacki, <i>Old Dominion University</i>	Glenn Ledder, <i>University of Nebraska—Lincoln</i>
Deborah Brandon, <i>Carnegie Mellon University</i>	Matthew Leingang, <i>New York University</i>
Samuel Chamberlin, <i>Park University</i>	Xin Li, <i>University of Central Florida</i>
Leonard Chastofsky, <i>University of Georgia</i>	Abey Lopez-Garcia, <i>University of South Alabama</i>
Meighan Dillon, <i>Southern Polytechnic State University</i>	Maura Mast, <i>University of Massachusetts—Boston</i>
Anne Dougherty, <i>University of Colorado</i>	Val Mohanakumar, <i>Hillsborough Community College—Dale Mabry Campus</i>
Said Fariabi, <i>San Antonio College</i>	Aaron Montgomery, <i>Central Washington University</i>
Klaus Fischer, <i>George Mason University</i>	Yibiao Pan, <i>University of Pittsburgh</i>
Tim Flood, <i>Pittsburg State University</i>	Christopher M. Pavone, <i>California State University at Chico</i>
Rick Ford, <i>California State University—Chico</i>	Cynthia Piez, <i>University of Idaho</i>
Toni Fountain, <i>Chattanooga State Community College</i>	Brooke Quinlan, <i>Hillsborough Community College—Dale Mabry Campus</i>
Robert Gardner, <i>East Tennessee State University</i>	Paul Sacks, <i>Iowa State University</i>
Mark Greer, <i>University of North Alabama</i>	Rebecca A. Segal, <i>Virginia Commonwealth University</i>
Ivan Gotchev, <i>Central Connecticut State University</i>	Andrew V. Sills, <i>Georgia Southern University</i>
Christopher Heil, <i>Georgia Institute of Technology</i>	Edward E. Slaminka, <i>Auburn University</i>
David Hemmer, <i>SUNY—Buffalo</i>	Alex Smith, <i>University of Wisconsin—Eau Claire</i>
Joshua Brandon Holden, <i>Rose-Hulman Institute of Technology</i>	Mark A. Smith, <i>Miami University</i>
Alexander Hulpke, <i>Colorado State University</i>	Donald Solomon, <i>University of Wisconsin—Milwaukee</i>
Jacqueline Jensen, <i>Sam Houston State University</i>	John Sullivan, <i>Black Hawk College</i>
Jennifer M. Johnson, <i>Princeton University</i>	Stephen Summers, <i>University of Florida</i>
	Maria Terrell, <i>Cornell University</i>

Blake Thornton, *Washington University
in St. Louis*

Ruth Trubnik, *Delaware Valley College*

Ilie Ugarcovici, *Rice University*

David Walnut, *George Mason University*

Adrian Wilson, *University of Montevallo*

Bobby Winters, *Pittsburg State
University*

Dennis Wortman, *University of
Massachusetts—Boston*

Yilian Zhang, *University of South
Carolina, Aiken*

The publishers would like to thank the following for their contribution to the Global Edition:

Contributor

Nalinakshi N., *Atria Institute of Technology*

Reviewers

B. R. Shankar, *National Institute of Technology Karnataka*

Antonio Behn Von Schmieden, *University of Chile*

Sunil Jacob John, *National Institute of Technology Calicut*

Credits

Chapter 3

Page 159, Exercise 19, PSSC PHYSICS, 2nd ed., Reprinted by permission of Education Development Center, Inc.

Chapter 6

Page 397, Figure 6.40, PSSC PHYSICS, 2nd ed., Reprinted by permission of Educational Development Center, Inc.

Chapter 9

Page 507, Figure 9.9, PSSC PHYSICS, 2nd ed., Reprinted by permission of Educational Development Center, Inc.

Chapter 13

Page 694, Figure 13.7, Reprinted by permission of the Appalachian Mountain Club;
Page 727, Figure 13.26, U.S. Geological Survey

Chapter 15

Page 842, Figures 15.6 and 15.7, National Committee for Fluid Mechanics, edited by Shapiro, ILLUSTRATED EXPERIMENTS IN FLUID MECHANICS: THE NCFMF BOOK OF FILM NOTES, reprinted by permission of Educational Development Center, Inc.; **Page 844, Figure 15.15**, InterNetwork Media, Inc., courtesy of NASA/JPL

Functions

OVERVIEW Functions are fundamental to the study of calculus. In this chapter we review what functions are and how they are pictured as graphs, how they are combined and transformed, and ways they can be classified. We review the trigonometric functions, and we discuss misrepresentations that can occur when using calculators and computers to obtain a function's graph. We also discuss inverse, exponential, and logarithmic functions. The real number system, Cartesian coordinates, straight lines, circles, parabolas, ellipses, and hyperbolas are reviewed in the Appendices.

1.1 Functions and Their Graphs

Functions are a tool for describing the real world in mathematical terms. A function can be represented by an equation, a graph, a numerical table, or a verbal description; we will use all four representations throughout this book. This section reviews these function ideas.

Functions; Domain and Range

The temperature at which water boils depends on the elevation above sea level (the boiling point drops as you ascend). The interest paid on a cash investment depends on the length of time the investment is held. The area of a circle depends on the radius of the circle. The distance an object travels at constant speed along a straight-line path depends on the elapsed time.

In each case, the value of one variable quantity, say y , depends on the value of another variable quantity, which we might call x . We say that “ y is a function of x ” and write this symbolically as

$$y = f(x) \quad (\text{“}y \text{ equals } f \text{ of } x\text{”}).$$

In this notation, the symbol f represents the function, the letter x is the **independent variable** representing the input value of f , and y is the **dependent variable** or output value of f at x .

DEFINITION A **function** f from a set D to a set Y is a rule that assigns a *unique* (single) element $f(x) \in Y$ to each element $x \in D$.

The set D of all possible input values is called the **domain** of the function. The set of all output values of $f(x)$ as x varies throughout D is called the **range** of the function. The range may not include every element in the set Y . The domain and range of a function can be any sets of objects, but often in calculus they are sets of real numbers interpreted as points of a coordinate line. (In Chapters 12–15, we will encounter functions for which the elements of the sets are points in the coordinate plane or in space.)




FIGURE 1.1 A diagram showing a function as a kind of machine.




FIGURE 1.2 A function from a set D to a set Y assigns a unique element of Y to each element in D .

Often a function is given by a formula that describes how to calculate the output value from the input variable. For instance, the equation $A = \pi r^2$ is a rule that calculates the area A of a circle from its radius r (so r , interpreted as a length, can only be positive in this formula). When we define a function $y = f(x)$ with a formula and the domain is not stated explicitly or restricted by context, the domain is assumed to be the largest set of real x -values for which the formula gives real y -values, which is called the **natural domain**. If we want to restrict the domain in some way, we must say so. The domain of $y = x^2$ is the entire set of real numbers. To restrict the domain of the function to, say, positive values of x , we would write “ $y = x^2, x > 0$.”

Changing the domain to which we apply a formula usually changes the range as well. The range of $y = x^2$ is $[0, \infty)$. The range of $y = x^2, x \geq 2$, is the set of all numbers obtained by squaring numbers greater than or equal to 2. In set notation (see Appendix 1), the range is $\{x^2 \mid x \geq 2\}$ or $\{y \mid y \geq 4\}$ or $[4, \infty)$.

When the range of a function is a set of real numbers, the function is said to be **real-valued**. The domains and ranges of most real-valued functions of a real variable we consider are intervals or combinations of intervals. The intervals may be open, closed, or half open, and may be finite or infinite. Sometimes the range of a function is not easy to find.

A function f is like a machine that produces an output value $f(x)$ in its range whenever we feed it an input value x from its domain (Figure 1.1). The function keys on a calculator give an example of a function as a machine. For instance, the \sqrt{x} key on a calculator gives an output value (the square root) whenever you enter a nonnegative number x and press the \sqrt{x} key.

A function can also be pictured as an **arrow diagram** (Figure 1.2). Each arrow associates an element of the domain D with a unique or single element in the set Y . In Figure 1.2, the arrows indicate that $f(a)$ is associated with a , $f(x)$ is associated with x , and so on. Notice that a function can have the same *value* at two different input elements in the domain (as occurs with $f(a)$ in Figure 1.2), but each input element x is assigned a *single* output value $f(x)$.

EXAMPLE 1 Let’s verify the natural domains and associated ranges of some simple functions. The domains in each case are the values of x for which the formula makes sense.

Function	Domain (x)	Range (y)
$y = x^2$	$(-\infty, \infty)$	$[0, \infty)$
$y = 1/x$	$(-\infty, 0) \cup (0, \infty)$	$(-\infty, 0) \cup (0, \infty)$
$y = \sqrt{x}$	$[0, \infty)$	$[0, \infty)$
$y = \sqrt{4 - x}$	$(-\infty, 4]$	$[0, \infty)$
$y = \sqrt{1 - x^2}$	$[-1, 1]$	$[0, 1]$

Solution The formula $y = x^2$ gives a real y -value for any real number x , so the domain is $(-\infty, \infty)$. The range of $y = x^2$ is $[0, \infty)$ because the square of any real number is nonnegative and every nonnegative number y is the square of its own square root, $y = (\sqrt{y})^2$ for $y \geq 0$.

The formula $y = 1/x$ gives a real y -value for every x except $x = 0$. For consistency in the rules of arithmetic, *we cannot divide any number by zero*. The range of $y = 1/x$, the set of reciprocals of all nonzero real numbers, is the set of all nonzero real numbers, since $y = 1/(1/y)$. That is, for $y \neq 0$ the number $x = 1/y$ is the input assigned to the output value y .

The formula $y = \sqrt{x}$ gives a real y -value only if $x \geq 0$. The range of $y = \sqrt{x}$ is $[0, \infty)$ because every nonnegative number is some number’s square root (namely, it is the square root of its own square).

In $y = \sqrt{4 - x}$, the quantity $4 - x$ cannot be negative. That is, $4 - x \geq 0$, or $x \leq 4$. The formula gives real y -values for all $x \leq 4$. The range of $\sqrt{4 - x}$ is $[0, \infty)$, the set of all nonnegative numbers.

The formula $y = \sqrt{1 - x^2}$ gives a real y -value for every x in the closed interval from -1 to 1 . Outside this domain, $1 - x^2$ is negative and its square root is not a real number. The values of $1 - x^2$ vary from 0 to 1 on the given domain, and the square roots of these values do the same. The range of $\sqrt{1 - x^2}$ is $[0, 1]$. ■

Graphs of Functions

If f is a function with domain D , its **graph** consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f . In set notation, the graph is

$$\{(x, f(x)) \mid x \in D\}.$$

The graph of the function $f(x) = x + 2$ is the set of points with coordinates (x, y) for which $y = x + 2$. Its graph is the straight line sketched in Figure 1.3.

The graph of a function f is a useful picture of its behavior. If (x, y) is a point on the graph, then $y = f(x)$ is the height of the graph above (or below) the point x . The height may be positive or negative, depending on the sign of $f(x)$ (Figure 1.4).

x	$y = x^2$
-2	4
-1	1
0	0
1	1
$\frac{3}{2}$	$\frac{9}{4}$
2	4




FIGURE 1.3 The graph of $f(x) = x + 2$ is the set of points (x, y) for which y has the value $x + 2$.




FIGURE 1.4 If (x, y) lies on the graph of f , then the value $y = f(x)$ is the height of the graph above the point x (or below x if $f(x)$ is negative).

EXAMPLE 2 Graph the function $y = x^2$ over the interval $[-2, 2]$.

Solution Make a table of xy -pairs that satisfy the equation $y = x^2$. Plot the points (x, y) whose coordinates appear in the table, and draw a *smooth* curve (labeled with its equation) through the plotted points (see Figure 1.5). ■





FIGURE 1.5 Graph of the function in Example 2.

How do we know that the graph of $y = x^2$ doesn't look like one of these curves?



To find out, we could plot more points. But how would we then connect *them*? The basic question still remains: How do we know for sure what the graph looks like between the points we plot? Calculus answers this question, as we will see in Chapter 4. Meanwhile, we will have to settle for plotting points and connecting them as best we can.

Representing a Function Numerically

We have seen how a function may be represented algebraically by a formula (the area function) and visually by a graph (Example 2). Another way to represent a function is **numerically**, through a table of values. Numerical representations are often used by engineers and experimental scientists. From an appropriate table of values, a graph of the function can be obtained using the method illustrated in Example 2, possibly with the aid of a computer. The graph consisting of only the points in the table is called a **scatterplot**.

EXAMPLE 3 Musical notes are pressure waves in the air. The data associated with Figure 1.6 give recorded pressure displacement versus time in seconds of a musical note produced by a tuning fork. The table provides a representation of the pressure function over time. If we first make a scatterplot and then connect approximately the data points (t, p) from the table, we obtain the graph shown in the figure.

Time	Pressure	Time	Pressure
0.00091	-0.080	0.00362	0.217
0.00108	0.200	0.00379	0.480
0.00125	0.480	0.00398	0.681
0.00144	0.693	0.00416	0.810
0.00162	0.816	0.00435	0.827
0.00180	0.844	0.00453	0.749
0.00198	0.771	0.00471	0.581
0.00216	0.603	0.00489	0.346
0.00234	0.368	0.00507	0.077
0.00253	0.099	0.00525	-0.164
0.00271	-0.141	0.00543	-0.320
0.00289	-0.309	0.00562	-0.354
0.00307	-0.348	0.00579	-0.248
0.00325	-0.248	0.00598	-0.035
0.00344	-0.041		




FIGURE 1.6 A smooth curve through the plotted points gives a graph of the pressure function represented by the accompanying tabled data (Example 3).

The Vertical Line Test for a Function

Not every curve in the coordinate plane can be the graph of a function. A function f can have only one value $f(x)$ for each x in its domain, so *no vertical* line can intersect the graph of a function more than once. If a is in the domain of the function f , then the vertical line $x = a$ will intersect the graph of f at the single point $(a, f(a))$.

A circle cannot be the graph of a function, since some vertical lines intersect the circle twice. The circle graphed in Figure 1.7a, however, does contain the graphs of functions of x , such as the upper semicircle defined by the function $f(x) = \sqrt{1 - x^2}$ and the lower semicircle defined by the function $g(x) = -\sqrt{1 - x^2}$ (Figures 1.7b and 1.7c).




FIGURE 1.7 (a) The circle is not the graph of a function; it fails the vertical line test. (b) The upper semicircle is the graph of a function $f(x) = \sqrt{1 - x^2}$. (c) The lower semicircle is the graph of a function $g(x) = -\sqrt{1 - x^2}$.




FIGURE 1.8 The absolute value function has domain $(-\infty, \infty)$ and range $[0, \infty)$.




FIGURE 1.9 To graph the function $y = f(x)$ shown here, we apply different formulas to different parts of its domain (Example 4).




FIGURE 1.10 The graph of the greatest integer function $y = \lfloor x \rfloor$ lies on or below the line $y = x$, so it provides an integer floor for x (Example 5).

Piecewise-Defined Functions

Sometimes a function is described in pieces by using different formulas on different parts of its domain. One example is the **absolute value function**

$$|x| = \begin{cases} x, & x \geq 0 & \text{First formula} \\ -x, & x < 0, & \text{Second formula} \end{cases}$$

whose graph is given in Figure 1.8. The right-hand side of the equation means that the function equals x if $x \geq 0$, and equals $-x$ if $x < 0$. Piecewise-defined functions often arise when real-world data are modeled. Here are some other examples.

EXAMPLE 4 The function

$$f(x) = \begin{cases} -x, & x < 0 & \text{First formula} \\ x^2, & 0 \leq x \leq 1 & \text{Second formula} \\ 1, & x > 1 & \text{Third formula} \end{cases}$$

is defined on the entire real line but has values given by different formulas, depending on the position of x . The values of f are given by $y = -x$ when $x < 0$, $y = x^2$ when $0 \leq x \leq 1$, and $y = 1$ when $x > 1$. The function, however, is *just one function* whose domain is the entire set of real numbers (Figure 1.9). ■

EXAMPLE 5 The function whose value at any number x is the *greatest integer less than or equal to* x is called the **greatest integer function** or the **integer floor function**. It is denoted $\lfloor x \rfloor$. Figure 1.10 shows the graph. Observe that

$$\begin{aligned} \lfloor 2.4 \rfloor &= 2, & \lfloor 1.9 \rfloor &= 1, & \lfloor 0 \rfloor &= 0, & \lfloor -1.2 \rfloor &= -2, \\ \lfloor 2 \rfloor &= 2, & \lfloor 0.2 \rfloor &= 0, & \lfloor -0.3 \rfloor &= -1, & \lfloor -2 \rfloor &= -2. \end{aligned}$$

EXAMPLE 6 The function whose value at any number x is the *smallest integer greater than or equal to* x is called the **least integer function** or the **integer ceiling function**. It is denoted $\lceil x \rceil$. Figure 1.11 shows the graph. For positive values of x , this function might represent, for example, the cost of parking x hours in a parking lot that charges \$1 for each hour or part of an hour. ■




FIGURE 1.11 The graph of the least integer function $y = [x]$ lies on or above the line $y = x$, so it provides an integer ceiling for x (Example 6).

Increasing and Decreasing Functions

If the graph of a function *climbs* or *rises* as you move from left to right, we say that the function is *increasing*. If the graph *descends* or *falls* as you move from left to right, the function is *decreasing*.

DEFINITIONS Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. If $f(x_2) > f(x_1)$ whenever $x_1 < x_2$, then f is said to be **increasing** on I .
2. If $f(x_2) < f(x_1)$ whenever $x_1 < x_2$, then f is said to be **decreasing** on I .

It is important to realize that the definitions of increasing and decreasing functions must be satisfied for *every* pair of points x_1 and x_2 in I with $x_1 < x_2$. Because we use the inequality $<$ to compare the function values, instead of \leq , it is sometimes said that f is *strictly* increasing or decreasing on I . The interval I may be finite (also called bounded) or infinite (unbounded) and by definition never consists of a single point (Appendix 1).

EXAMPLE 7 The function graphed in Figure 1.9 is decreasing on $(-\infty, 0]$ and increasing on $[0, 1]$. The function is neither increasing nor decreasing on the interval $[1, \infty)$ because of the strict inequalities used to compare the function values in the definitions. ■

Even Functions and Odd Functions: Symmetry

The graphs of *even* and *odd* functions have characteristic symmetry properties.

DEFINITIONS A function $y = f(x)$ is an

even function of x if $f(-x) = f(x)$,

odd function of x if $f(-x) = -f(x)$,

for every x in the function's domain.




FIGURE 1.12 (a) The graph of $y = x^2$ (an even function) is symmetric about the y -axis. (b) The graph of $y = x^3$ (an odd function) is symmetric about the origin.

The names *even* and *odd* come from powers of x . If y is an even power of x , as in $y = x^2$ or $y = x^4$, it is an even function of x because $(-x)^2 = x^2$ and $(-x)^4 = x^4$. If y is an odd power of x , as in $y = x$ or $y = x^3$, it is an odd function of x because $(-x)^1 = -x$ and $(-x)^3 = -x^3$.

The graph of an even function is **symmetric about the y -axis**. Since $f(-x) = f(x)$, a point (x, y) lies on the graph if and only if the point $(-x, y)$ lies on the graph (Figure 1.12a). A reflection across the y -axis leaves the graph unchanged.

The graph of an odd function is **symmetric about the origin**. Since $f(-x) = -f(x)$, a point (x, y) lies on the graph if and only if the point $(-x, -y)$ lies on the graph (Figure 1.12b). Equivalently, a graph is symmetric about the origin if a rotation of 180° about the origin leaves the graph unchanged. Notice that the definitions imply that both x and $-x$ must be in the domain of f .

EXAMPLE 8 Here are several functions illustrating the definition.

$f(x) = x^2$ Even function: $(-x)^2 = x^2$ for all x ; symmetry about y -axis.

$f(x) = x^2 + 1$ Even function: $(-x)^2 + 1 = x^2 + 1$ for all x ; symmetry about y -axis (Figure 1.13a).

$f(x) = x$ Odd function: $(-x) = -x$ for all x ; symmetry about the origin.

$f(x) = x + 1$ Not odd: $f(-x) = -x + 1$, but $-f(x) = -x - 1$. The two are not equal.

Not even: $(-x) + 1 \neq x + 1$ for all $x \neq 0$ (Figure 1.13b). ■




FIGURE 1.13 (a) When we add the constant term 1 to the function $y = x^2$, the resulting function $y = x^2 + 1$ is still even and its graph is still symmetric about the y -axis. (b) When we add the constant term 1 to the function $y = x$, the resulting function $y = x + 1$ is no longer odd, since the symmetry about the origin is lost. The function $y = x + 1$ is also not even (Example 8).

Common Functions

A variety of important types of functions are frequently encountered in calculus. We identify and briefly describe them here.

Linear Functions A function of the form $f(x) = mx + b$, for constants m and b , is called a **linear function**. Figure 1.14a shows an array of lines $f(x) = mx$ where $b = 0$, so these lines pass through the origin. The function $f(x) = x$ where $m = 1$ and $b = 0$ is called the **identity function**. Constant functions result when the slope $m = 0$ (Figure 1.14b). A linear function with positive slope whose graph passes through the origin is called a *proportionality* relationship.




FIGURE 1.14 (a) Lines through the origin with slope m . (b) A constant function with slope $m = 0$.

DEFINITION Two variables y and x are **proportional** (to one another) if one is always a constant multiple of the other; that is, if $y = kx$ for some nonzero constant k .

If the variable y is proportional to the reciprocal $1/x$, then sometimes it is said that y is **inversely proportional** to x (because $1/x$ is the multiplicative inverse of x).

Power Functions A function $f(x) = x^a$, where a is a constant, is called a **power function**. There are several important cases to consider.

(a) $a = n$, a positive integer.

The graphs of $f(x) = x^n$, for $n = 1, 2, 3, 4, 5$, are displayed in Figure 1.15. These functions are defined for all real values of x . Notice that as the power n gets larger, the curves tend to flatten toward the x -axis on the interval $(-1, 1)$, and to rise more steeply for $|x| > 1$. Each curve passes through the point $(1, 1)$ and through the origin. The graphs of functions with even powers are symmetric about the y -axis; those with odd powers are symmetric about the origin. The even-powered functions are decreasing on the interval $(-\infty, 0]$ and increasing on $[0, \infty)$; the odd-powered functions are increasing over the entire real line $(-\infty, \infty)$.




FIGURE 1.15 Graphs of $f(x) = x^n$, $n = 1, 2, 3, 4, 5$, defined for $-\infty < x < \infty$.

(b) $a = -1$ or $a = -2$.

The graphs of the functions $f(x) = x^{-1} = 1/x$ and $g(x) = x^{-2} = 1/x^2$ are shown in Figure 1.16. Both functions are defined for all $x \neq 0$ (you can never divide by zero). The graph of $y = 1/x$ is the hyperbola $xy = 1$, which approaches the coordinate axes far from the origin. The graph of $y = 1/x^2$ also approaches the coordinate axes. The graph of the function f is symmetric about the origin; f is decreasing on the intervals $(-\infty, 0)$ and $(0, \infty)$. The graph of the function g is symmetric about the y -axis; g is increasing on $(-\infty, 0)$ and decreasing on $(0, \infty)$.




FIGURE 1.16 Graphs of the power functions $f(x) = x^a$ for part (a) $a = -1$ and for part (b) $a = -2$.

(c) $a = \frac{1}{2}, \frac{1}{3}, \frac{3}{2}$, and $\frac{2}{3}$.

The functions $f(x) = x^{1/2} = \sqrt{x}$ and $g(x) = x^{1/3} = \sqrt[3]{x}$ are the **square root** and **cube root** functions, respectively. The domain of the square root function is $[0, \infty)$, but the cube root function is defined for all real x . Their graphs are displayed in Figure 1.17, along with the graphs of $y = x^{3/2}$ and $y = x^{2/3}$. (Recall that $x^{3/2} = (x^{1/2})^3$ and $x^{2/3} = (x^{1/3})^2$.)

Polynomials A function p is a **polynomial** if

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

where n is a nonnegative integer and the numbers $a_0, a_1, a_2, \dots, a_n$ are real constants (called the **coefficients** of the polynomial). All polynomials have domain $(-\infty, \infty)$. If the




FIGURE 1.17 Graphs of the power functions $f(x) = x^a$ for $a = \frac{1}{2}$, $\frac{1}{3}$, $\frac{3}{2}$, and $\frac{2}{3}$.

leading coefficient $a_n \neq 0$ and $n > 0$, then n is called the **degree** of the polynomial. Linear functions with $m \neq 0$ are polynomials of degree 1. Polynomials of degree 2, usually written as $p(x) = ax^2 + bx + c$, are called **quadratic functions**. Likewise, **cubic functions** are polynomials $p(x) = ax^3 + bx^2 + cx + d$ of degree 3. Figure 1.18 shows the graphs of three polynomials. Techniques to graph polynomials are studied in Chapter 4.




FIGURE 1.18 Graphs of three polynomial functions.

Rational Functions A **rational function** is a quotient or ratio $f(x) = p(x)/q(x)$, where p and q are polynomials. The domain of a rational function is the set of all real x for which $q(x) \neq 0$. The graphs of several rational functions are shown in Figure 1.19.




FIGURE 1.19 Graphs of three rational functions. The straight red lines approached by the graphs are called *asymptotes* and are not part of the graphs. We discuss asymptotes in Section 2.6.

Algebraic Functions Any function constructed from polynomials using algebraic operations (addition, subtraction, multiplication, division, and taking roots) lies within the class of **algebraic functions**. All rational functions are algebraic, but also included are more complicated functions (such as those satisfying an equation like $y^3 - 9xy + x^3 = 0$, studied in Section 3.7). Figure 1.20 displays the graphs of three algebraic functions.




FIGURE 1.20 Graphs of three algebraic functions.

Trigonometric Functions The six basic trigonometric functions are reviewed in Section 1.3. The graphs of the sine and cosine functions are shown in Figure 1.21.




FIGURE 1.21 Graphs of the sine and cosine functions.

Exponential Functions Functions of the form $f(x) = a^x$, where the base $a > 0$ is a positive constant and $a \neq 1$, are called **exponential functions**. All exponential functions have domain $(-\infty, \infty)$ and range $(0, \infty)$, so an exponential function never assumes the value 0. We discuss exponential functions in Section 1.5. The graphs of some exponential functions are shown in Figure 1.22.




FIGURE 1.22 Graphs of exponential functions.

Logarithmic Functions These are the functions $f(x) = \log_a x$, where the base $a \neq 1$ is a positive constant. They are the *inverse functions* of the exponential functions, and we discuss these functions in Section 1.6. Figure 1.23 shows the graphs of four logarithmic functions with various bases. In each case the domain is $(0, \infty)$ and the range is $(-\infty, \infty)$.




FIGURE 1.23 Graphs of four logarithmic functions.




FIGURE 1.24 Graph of a catenary or hanging cable. (The Latin word *catena* means “chain.”)

Transcendental Functions These are functions that are not algebraic. They include the trigonometric, inverse trigonometric, exponential, and logarithmic functions, and many other functions as well. A particular example of a transcendental function is a **catenary**. Its graph has the shape of a cable, like a telephone line or electric cable, strung from one support to another and hanging freely under its own weight (Figure 1.24). The function defining the graph is discussed in Section 7.3.



Exercises 1.1

Functions

In Exercises 1–6, find the domain and range of each function.

- $f(x) = 18 + x^2$
- $f(x) = 1 - \sqrt{x}$
- $F(x) = \sqrt{5x + 10}$
- $g(x) = \sqrt{x^2 - 3x}$
- $f(t) = \frac{4}{3 - t}$
- $G(t) = \frac{5}{t^2 - 4}$

In Exercises 7 and 8, which of the graphs are graphs of functions of x , and which are not? Give reasons for your answers.



Finding Formulas for Functions

- Express the area and perimeter of an equilateral triangle as a function of the triangle's side length x .
- Express the side length of a square as a function of the length d of the square's diagonal. Then express the area as a function of the diagonal length.
- Express the edge length of a cube as a function of the cube's diagonal length d . Then express the surface area and volume of the cube as a function of the diagonal length.

12. A point P in the first quadrant lies on the graph of the function $f(x) = \sqrt{x}$. Express the coordinates of P as functions of the slope of the line joining P to the origin.
13. Consider the point (x, y) lying on the graph of the line $2x + 4y = 5$. Let L be the distance from the point (x, y) to the origin $(0, 0)$. Write L as a function of x .
14. Consider the point (x, y) lying on the graph of $y = \sqrt{x - 3}$. Let L be the distance between the points (x, y) and $(4, 0)$. Write L as a function of y .

Functions and Graphs

Find the natural domain and graph the functions in Exercises 15–20.

15. $f(x) = 5 - 2x$ 16. $f(x) = 1 - 2x - x^2$
 17. $g(x) = \sqrt{|x|}$ 18. $g(x) = \sqrt{-x}$
 19. $F(t) = t/|t|$ 20. $G(t) = 1/|t|$

21. Find the domain of $y = \frac{x + 7}{12 - \sqrt{x^2 - 25}}$.
22. Find the range of $y = 2 + \frac{x^2}{x^2 + 4}$.
23. Graph the following equations and explain why they are not graphs of functions of x .
 a. $|y| = x$ b. $y^2 = x^2$
24. Graph the following equations and explain why they are not graphs of functions of x .
 a. $|x| + |y| = 1$ b. $|x + y| = 1$

Piecewise-Defined Functions

Graph the functions in Exercises 25–28.

25. $f(x) = \begin{cases} x, & 0 \leq x \leq 1 \\ 2 - x, & 1 < x \leq 2 \end{cases}$
 26. $g(x) = \begin{cases} 1 - x, & 0 \leq x \leq 1 \\ 2 - x, & 1 < x \leq 2 \end{cases}$
 27. $F(x) = \begin{cases} 4 - x^2, & x \leq 1 \\ x^2 + 2x, & x > 1 \end{cases}$
 28. $G(x) = \begin{cases} 1/x, & x < 0 \\ x, & 0 \leq x \end{cases}$

Find a formula for each function graphed in Exercises 29–32.

29. a. b.
30. a. b.

31. a. b.
32. a. b.

The Greatest and Least Integer Functions

33. For what values of x is
 a. $\lfloor x \rfloor = -1$? b. $\lceil x \rceil = 0$?
34. What real numbers x satisfy the equation $\lfloor x \rfloor = \lceil x \rceil$?
35. Does $\lceil -x \rceil = -\lfloor x \rfloor$ for all real x ? Give reasons for your answer.
36. Graph the function

$$f(x) = \begin{cases} \lfloor x \rfloor, & x \geq 0 \\ \lceil x \rceil, & x < 0. \end{cases}$$

Why is $f(x)$ called the *integer part* of x ?

Increasing and Decreasing Functions

Graph the functions in Exercises 37–46. What symmetries, if any, do the graphs have? Specify the intervals over which the function is increasing and the intervals where it is decreasing.

37. $y = -x^3$ 38. $y = -\frac{1}{x^2}$
 39. $y = -\frac{1}{x}$ 40. $y = \frac{1}{|x|}$
 41. $y = \sqrt{|x|}$ 42. $y = \sqrt{-x}$
 43. $y = x^3/8$ 44. $y = -4\sqrt{x}$
 45. $y = -x^{3/2}$ 46. $y = (-x)^{2/3}$

Even and Odd Functions


In Exercises 47–58, say whether the function is even, odd, or neither. Give reasons for your answer.

47. $f(x) = -8$ 48. $f(x) = x^{-5}$
 49. $g(x) = 9x^3 - 3$ 50. $f(x) = x^2 + x$
 51. $g(x) = x^3 + x$ 52. $g(x) = x^4 + 3x^2 - 1$
 53. $g(x) = \frac{1}{x^2 - 4}$ 54. $g(x) = \frac{x}{x^2 - 1}$
 55. $h(t) = \frac{1}{t^3 + 1}$ 56. $h(t) = |t^3|$
 57. $h(t) = 5t - 1$ 58. $h(t) = 2|t| + 1$


Theory and Examples

59. The variable s is proportional to t , and $s = 15$ when $t = 105$. Determine t when $s = 40$.

60. **Kinetic energy** The kinetic energy K of a mass is proportional to the square of its velocity v . If $K = 12,960$ joules when $v = 18$ m/sec, what is K when $v = 10$ m/sec?
61. The variables r and s are inversely proportional, and $r = 7$ when $s = 4$. Determine s when $r = 10$.
62. **Boyle's Law** Boyle's Law says that the volume V of a gas at constant temperature increases whenever the pressure P decreases, so that V and P are inversely proportional. If $P = 14.7$ lb/in² when $V = 1000$ in³, then what is V when $P = 23.4$ lb/in²?
63. A box with an open top is to be constructed from a rectangular piece of cardboard with dimensions 14 in. by 22 in. by cutting out equal squares of side x at each corner and then folding up the sides as in the figure. Express the volume V of the box as a function of x .




64. The accompanying figure shows a rectangle inscribed in an isosceles right triangle whose hypotenuse is 2 units long.
- Express the y -coordinate of P in terms of x . (You might start by writing an equation for the line AB .)
 - Express the area of the rectangle in terms of x .




In Exercises 65 and 66, match each equation with its graph. Do not use a graphing device, and give reasons for your answer.

65. a. $y = x^4$ b. $y = x^7$ c. $y = x^{10}$



66. a. $y = 5x$ b. $y = 5^x$ c. $y = x^5$



- T 67. a. Graph the functions $f(x) = x/2$ and $g(x) = 1 + (4/x)$ together to identify the values of x for which

$$\frac{x}{2} > 1 + \frac{4}{x}.$$


- b. Confirm your findings in part (a) algebraically.

- T 68. a. Graph the functions $f(x) = 3/(x - 1)$ and $g(x) = 2/(x + 1)$ together to identify the values of x for which

$$\frac{3}{x - 1} < \frac{2}{x + 1}.$$

- b. Confirm your findings in part (a) algebraically.

69. For a curve to be *symmetric about the x-axis*, the point (x, y) must lie on the curve if and only if the point $(x, -y)$ lies on the curve. Explain why a curve that is symmetric about the x -axis is not the graph of a function, unless the function is $y = 0$.
70. Three hundred books sell for \$40 each, resulting in a revenue of $(300)(\$40) = \$12,000$. For each \$5 increase in the price, 25 fewer books are sold. Write the revenue R as a function of the number x of \$5 increases.
71. A pen in the shape of an isosceles right triangle with legs of length x ft and hypotenuse of length h ft is to be built. If fencing costs \$2/ft for the legs and \$8/ft for the hypotenuse, write the total cost C of construction as a function of h .
72. **Industrial costs** A power plant sits next to a river where the river is 800 ft wide. To lay a new cable from the plant to a location in the city 2 mi downstream on the opposite side costs \$180 per foot across the river and \$100 per foot along the land.



- Suppose that the cable goes from the plant to a point Q on the opposite side that is x ft from the point P directly opposite the plant. Write a function $C(x)$ that gives the cost of laying the cable in terms of the distance x .
- Generate a table of values to determine if the least expensive location for point Q is less than 2000 ft or greater than 2000 ft from point P .

1.2 Combining Functions; Shifting and Scaling Graphs

In this section we look at the main ways functions are combined or transformed to form new functions.

Sums, Differences, Products, and Quotients

Like numbers, functions can be added, subtracted, multiplied, and divided (except where the denominator is zero) to produce new functions. If f and g are functions, then for every x that belongs to the domains of both f and g (that is, for $x \in D(f) \cap D(g)$), we define functions $f + g$, $f - g$, and fg by the formulas

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(fg)(x) = f(x)g(x).$$

Notice that the $+$ sign on the left-hand side of the first equation represents the operation of addition of *functions*, whereas the $+$ on the right-hand side of the equation means addition of the real numbers $f(x)$ and $g(x)$.

At any point of $D(f) \cap D(g)$ at which $g(x) \neq 0$, we can also define the function f/g by the formula

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad (\text{where } g(x) \neq 0).$$

Functions can also be multiplied by constants: If c is a real number, then the function cf is defined for all x in the domain of f by

$$(cf)(x) = cf(x).$$

EXAMPLE 1 The functions defined by the formulas

$$f(x) = \sqrt{x} \quad \text{and} \quad g(x) = \sqrt{1-x}$$

have domains $D(f) = [0, \infty)$ and $D(g) = (-\infty, 1]$. The points common to these domains are the points

$$[0, \infty) \cap (-\infty, 1] = [0, 1].$$

The following table summarizes the formulas and domains for the various algebraic combinations of the two functions. We also write $f \cdot g$ for the product function fg .

Function	Formula	Domain
$f + g$	$(f + g)(x) = \sqrt{x} + \sqrt{1-x}$	$[0, 1] = D(f) \cap D(g)$
$f - g$	$(f - g)(x) = \sqrt{x} - \sqrt{1-x}$	$[0, 1]$
$g - f$	$(g - f)(x) = \sqrt{1-x} - \sqrt{x}$	$[0, 1]$
$f \cdot g$	$(f \cdot g)(x) = f(x)g(x) = \sqrt{x(1-x)}$	$[0, 1]$
f/g	$\frac{f}{g}(x) = \frac{f(x)}{g(x)} = \sqrt{\frac{x}{1-x}}$	$[0, 1)$ ($x = 1$ excluded)
g/f	$\frac{g}{f}(x) = \frac{g(x)}{f(x)} = \sqrt{\frac{1-x}{x}}$	$(0, 1]$ ($x = 0$ excluded)

The graph of the function $f + g$ is obtained from the graphs of f and g by adding the corresponding y -coordinates $f(x)$ and $g(x)$ at each point $x \in D(f) \cap D(g)$, as in Figure 1.25. The graphs of $f + g$ and $f \cdot g$ from Example 1 are shown in Figure 1.26.




FIGURE 1.25 Graphical addition of two functions.




FIGURE 1.26 The domain of the function $f + g$ is the intersection of the domains of f and g , the interval $[0, 1]$ on the x -axis where these domains overlap. This interval is also the domain of the function $f \cdot g$ (Example 1).

Composite Functions

Composition is another method for combining functions.

DEFINITION If f and g are functions, the **composite** function $f \circ g$ (“ f composed with g ”) is defined by

$$(f \circ g)(x) = f(g(x)).$$

The domain of $f \circ g$ consists of the numbers x in the domain of g for which $g(x)$ lies in the domain of f .

The definition implies that $f \circ g$ can be formed when the range of g lies in the domain of f . To find $(f \circ g)(x)$, *first* find $g(x)$ and *second* find $f(g(x))$. Figure 1.27 pictures $f \circ g$ as a machine diagram, and Figure 1.28 shows the composite as an arrow diagram.




FIGURE 1.27 A composite function $f \circ g$ uses the output $g(x)$ of the first function g as the input for the second function f .




FIGURE 1.28 Arrow diagram for $f \circ g$. If x lies in the domain of g and $g(x)$ lies in the domain of f , then the functions f and g can be composed to form $(f \circ g)(x)$.

To evaluate the composite function $g \circ f$ (when defined), we find $f(x)$ first and then $g(f(x))$. The domain of $g \circ f$ is the set of numbers x in the domain of f such that $f(x)$ lies in the domain of g .

The functions $f \circ g$ and $g \circ f$ are usually quite different.

EXAMPLE 2 If $f(x) = \sqrt{x}$ and $g(x) = x + 1$, find

- (a) $(f \circ g)(x)$ (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$.

Solution

Composite	Domain
(a) $(f \circ g)(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x + 1}$	$[-1, \infty)$
(b) $(g \circ f)(x) = g(f(x)) = f(x) + 1 = \sqrt{x} + 1$	$[0, \infty)$
(c) $(f \circ f)(x) = f(f(x)) = \sqrt{f(x)} = \sqrt{\sqrt{x}} = x^{1/4}$	$[0, \infty)$
(d) $(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x + 1) + 1 = x + 2$	$(-\infty, \infty)$

To see why the domain of $f \circ g$ is $[-1, \infty)$, notice that $g(x) = x + 1$ is defined for all real x but belongs to the domain of f only if $x + 1 \geq 0$, that is to say, when $x \geq -1$. ■

Notice that if $f(x) = x^2$ and $g(x) = \sqrt{x}$, then $(f \circ g)(x) = (\sqrt{x})^2 = x$. However, the domain of $f \circ g$ is $[0, \infty)$, not $(-\infty, \infty)$, since \sqrt{x} requires $x \geq 0$.

Shifting a Graph of a Function

A common way to obtain a new function from an existing one is by adding a constant to each output of the existing function, or to its input variable. The graph of the new function is the graph of the original function shifted vertically or horizontally, as follows.

Shift Formulas

Vertical Shifts

$y = f(x) + k$ Shifts the graph of f up k units if $k > 0$
 Shifts it down $|k|$ units if $k < 0$

Horizontal Shifts

$y = f(x + h)$ Shifts the graph of f left h units if $h > 0$
 Shifts it right $|h|$ units if $h < 0$




FIGURE 1.29 To shift the graph of $f(x) = x^2$ up (or down), we add positive (or negative) constants to the formula for f (Examples 3a and b).

EXAMPLE 3

- (a) Adding 1 to the right-hand side of the formula $y = x^2$ to get $y = x^2 + 1$ shifts the graph up 1 unit (Figure 1.29).
 (b) Adding -2 to the right-hand side of the formula $y = x^2$ to get $y = x^2 - 2$ shifts the graph down 2 units (Figure 1.29).
 (c) Adding 3 to x in $y = x^2$ to get $y = (x + 3)^2$ shifts the graph 3 units to the left, while adding -2 shifts the graph 2 units to the right (Figure 1.30).
 (d) Adding -2 to x in $y = |x|$, and then adding -1 to the result, gives $y = |x - 2| - 1$ and shifts the graph 2 units to the right and 1 unit down (Figure 1.31). ■

Scaling and Reflecting a Graph of a Function

To scale the graph of a function $y = f(x)$ is to stretch or compress it, vertically or horizontally. This is accomplished by multiplying the function f , or the independent variable x , by an appropriate constant c . Reflections across the coordinate axes are special cases where $c = -1$.